



Review Article

www.ijrap.net

(ISSN Online:2229-3566, ISSN Print:2277-4343)



STHAULYA, AMA, AND MEDA DHATU, THE DYSFUNCTIONAL NUTRITIONAL METABOLISM WITH SPECIAL REFERENCE TO OXIDATIVE STRESS: A NARRATIVE REVIEW

Amulya Murthy Aku ^{1*}, Ashok Patil ², Mukund Sabnis ³

¹ PG Scholar, Department of PG Studies in Swasthavritta, K.L.E. Academy of Higher Education & Research's (K.A.H.E.R., Deemed-to-be-University) Shri B. M. Kankanawadi Ayurveda Mahavidyalaya, Shahpur, Belgaum, Karnataka, India

² Professor & HOD, Department of PG Studies in Swasthavritta, K.L.E. Academy of Higher Education & Research's (K.A.H.E.R., Deemed-to-be-University) Shri B. M. Kankanawadi Ayurveda Mahavidyalaya, Shahpur, Belgaum, Karnataka, India

³ Specialist Clinician, Founder President of Jeevan Rekha Ayurveda Chikitsalaya & Research Centre, Jawahar Colony, Aurangabad, Maharashtra, India

Received on: 02/02/22 Accepted on: 26/04/22

***Corresponding author**

E-mail: amulyamurthy2016@gmail.com

DOI: 10.7897/2277-4343.130496

ABSTRACT

Obesity and its accompanying consequences are increasingly linked to oxidative stress, as evidenced by a growing body of research. Several biochemical pathways, including superoxide production from NADPH oxidases and glyceraldehyde auto-oxidation, can cause systemic oxidative stress in obese individuals. Hyperleptinemia, inadequate antioxidant defence, chronic inflammation, and postprandial reactive oxygen species production contribute to oxidative stress. This Oxidative stress is related to or co-related to "Ama". Ama is generated by improper or insufficient digestion of food particles due to a lack of *Jatharagni* and *Mala* accumulation in the body. Modern physiology has linked Ama to potentially harmful Free radicals. It is essential to neutralize free radicals because they have one or more unpaired electrons. Oxygen-free radicals, or "Reactive Oxygen Species," make up most dangerous free radicals. Ama is in the same boat as *Avipakam* (incompletely metabolized) due to its metabolism. This Ama brings on many diseases. Many diseases have been linked to free radicals.

Keywords: *Sthaulya*, Obesity, Oxidative stress, *Ama*, Free Radicals, NADPH

INTRODUCTION

Obesity is a chronic disease that affects people's physical, financial, and mental well-being, regardless of their cultural, economic, or ethnic background. Excess body fat lowers the quality of life and increases healthcare costs and death risk. Diabetes, cardiovascular disease, cancer, asthma, sleep disorders, hepatic, renal, and infertility are all associated with obesity.¹⁻⁴ Overweight is defined as a BMI of 25.0 to 29.9 kg/m², while obesity is defined as a BMI of 30 kg/m². BMI,⁵ on the other hand, does not distinguish between lean and fat mass, nor does it specify fat distribution as a distinguishing factor. Obesity-related risk factors are found to be less dependent on total body weight and more dependent on fat distribution, according to new research.⁶⁻⁷ As a result, because it promotes pro-oxidant and pro-inflammatory states, abdominal fat is now recognized as a significant risk factor for obesity-related illnesses.⁸⁻¹⁰

This ailment is covered under *Sthaulya* in *Ayurveda*, the science of life. Because of its complicated pathologies and multifactor aetiologies, *Sthaulya* is expected to be extremely difficult to treat. The creation of *Ama* is the fundamental cause of these complexities. Because of the long-term ingestion of *Nidaana* (aetiologies), the body cannot clear the toxins created because of these aetiologies, and these toxins accumulate in the form of *Ama*. This *Ama* produces a layer all over the *Jathargni*, interfering with its function, and creates *Saama Dhatus* by interfering with *Dhatuagni's* function. This causes an excessive accumulation of

Meda Dhatu, resulting in *Sthaulya* illness. This *Ama* is frequently linked to Free Radicals, ROS, that cause oxidative stress.

Oxidative stress has been associated with obesity in epidemiological, clinical, and animal studies. In cell culture and animal studies, oxidative stress increases pre-adipocyte proliferation, differentiation, and mature adipocyte size.¹¹ Reactive oxygen species have been proven to impact hypothalamic satiety and hunger neurons (ROS). N-acetyl-glycer-aldehyde auto-oxidation, polyol and hexose-amine routes are all produced by NOXPH oxidases. Obesity causes oxidative stress due to poor antioxidant defence, chronic inflammation, and postprandial ROS generation.¹²⁻¹⁶ This paper attempts to cover the role of *Ama* in low-quality *Dhatu* formation, focusing on *Meda Dhatu*.

Literature

Literature on *Ama* (Free radicals), *Sthaulya* (Obesity), and *Meda Dhatu Poshana* (Adipose tissue nourishment and replenishment) was gathered from various *Ayurvedic* Compendia. Current literature and recent research publications were also searched online from scientific sites such as PubMed, Google Scholar, and Medline plus using keywords such as *Ama*, Free Radicals in *Ayurveda*, *Meda dhatu*, and *Sthaulya*. Analysing the collected descriptions allowed researchers to determine the applicability and relevance of *Dhatu Poshana* dysfunction to *Sthaulya*.

DISCUSSION

Ayurveda's core principle is keeping *Dosha*, *Dhatu*, and *Mala* balanced. The potency of each *Jatharagni* initially nourishes these three, and then productive nutrients (*Ahara Rasa*) are passed into each level of *Dhatu* (body tissues). One pool provides all the nutrients required for tissue creation and growth.¹⁷ Among the three *Upasthambhas* (*Ahara*, *Nidra*, *Brahmacharya*), *Ahara* is the most important. Traditional texts explain that usually, the body follows three laws to fulfil the nutritional needs of the body by converting the food into such form so that it is absorbable (*Ksheera Dadhi Nyaya* – transformation of nutrients, *Khale Kapota Nyaya* - selective uptake of nutrients, *Kedari Kulya Nyaya* – transportation of nutrients via channels) but in obesity, such mechanism is deeply affected

***Kshira Dadhi Nyaya* (Metabolic transformation of nutrients)**

Kshira (milk) is distinct from *Dadhi* (curd). Milk undergoes a complete change before becoming curd. According to this hypothesis, a certain tissue undergoes differentiation into subsequent tissues. *Rasa Dhatu*, like milk, transforms into *Rakta Dhatu* following this law. Like *Mamsa* and *Meda*, *Asthi* and *Majja* and *Shukra* are generated by the gradual stage of transformation, and *Shukra* is accountable for the cause of pregnancy. *Shukra* (garbha). *Karma Parinama Paksha* is another name for this.¹⁸⁻¹⁹

***Khale Kapota Nyaya* (Selective Uptake of Nutrients)**

After harvesting, the grain is stored in the *Khale* (granary or bran), and *Kapot* is a pigeon or bird symbol. For example, Pigeons visit a grain pile to pick up the requisite and return to their nest. Energy is used in this procedure by *Kapotas* to meet their needs. As with pigeons (*Kapotas*), the *Rasa Dhatu* contains the nutrient components for all the *Dhatu*s in the body, just like the *Rasa Dhatu* contains the grains for the *Dhatu*s.^{18,20,21}

***Kedari Kulya Nyaya* (Transportation of nutrients via channel)**

In this context, *Kulya* refers to a little waterway in a paddy field called *Kedari* (*Kyari*). Water from the reservoir is delivered to tiny fields (*Kedari*) via canals as part of the irrigation process. Water moves in *Kulya* (canals) in the direction of gravitational pull. The water reaches the first section of the field (*Kedari*) and then goes on to the second half of the field via a pressure gradient (considering the first *Kedari* is filled with water, then water moves to the next *Kedari*). This mechanism explains passive diffusion through cell membranes. *Meda Dhatu* and Obesity.^{18,20,22}

Dysfunction

Dysfunction of *Kshira Dadhi nayaya* in *Ama* and *Sthaulya*

In the case of *Ama* formation, the formed *Ahara Rasa* is coated with *Ama*, known as “*Sama Anna Rasa*”, because the subsequent *Dhatu*s made from it, i.e. *Rasa*, *Rakta*, *Mamsa*, *Meda Dhatu* is also covered with *Ama* and creates *Sama Dhatu*s.

Dysfunction of “*Khale Kapot Nayaya*” in *Ama* and *Sthaulya*

Similarly, the consecutive *Dhatu*s formed is of the lowest nutritional Quality and high in *Kapha*, *Guru*, *Snigdha* and similar properties. Therefore, the *Dhatu*s – *Rasa*, *Rakta*, *Mamsa*, and *Meda* cannot perform their optimum.

Dysfunction of “*Kedari Kulya Nyaya*” in *Ama* and *Sthaulya*

However, in this situation, the *Dhatu*s up to the point of *Meda* has been somewhat produced. Still, the nutrient pool is depleted, and

the next *Dhatu* cannot pick up essential attributes required for nourishment. The *Meda Dhatuagni* has been transformed into *Manda- Medadhatwagni*, preventing further transformation.

The mode of action and the probable reason behind this process needs to be understood at the level of *Rasa Dhatu* and then *Meda Dhatu*.

RASA DHATU CAN BE COMPARED TO PLASMA

Intracellular antioxidant defences must be adequate to maintain the balance of tissues' antioxidant-pro-oxidant. Vitamin and mineral deficiencies can also impair antioxidant defence in the pathogenesis of obesity. Plasma levels of α -tocopherol or β -carotene per unit of plasma low-density lipoprotein (LDL) are well-known antioxidant protection biomarkers. Low carotenoids, vitamin C, and vitamin E levels have increased BMI. The CARDIA study found an inverse connection between BMI and total serum carotenoids (α -carotene, β -carotene, α -cryptoxanthin, and zeaxanthin/lutein). Obese children had lower serum β -carotene levels than normal control children in the National Health and Examination Survey. In one study, obese girls had lower plasma levels of α -tocopherol/LDL and β -carotene/LDL than non-obese girls. According to Aasheim and Bohmer, most obese patients have low vitamin levels, particularly A, B6, C, D, and E. Dietary vitamin C intake was inversely associated with central fat distribution in the European Prospective Population Study (EPIC). People with BMI >50 had lower plasma levels of vitamin E/triglycerides than people with BMI 30. Obese children, especially those with central obesity, had more deficient selenium and zinc levels. Obese people also have magnesium, selenium, iron, and zinc deficiency. These findings imply that low vitamin and mineral concentrations in obese people promote reduced antioxidant defence.²³⁻³⁴

MEDA DHATU CAN BE CORRELATED TO WHITE ADIPOSE TISSUE

Plasma-free fatty acids (FFA) rise in obesity, as does fat storage in the waist and hips (WAT). The mitochondrial electron transport chain generates O₂ - by blocking the translocation of adenine nucleotides. At that point, the plasma FFA levels are elevated. FFA activates NOX via PKC in cultured vascular cells to produce reactive intermediates. As the conjugated fatty acids encounter oxygen, they become sensitive to oxidation, which causes the generation of free radicals and the accumulation of oxidative by-products. Fats are more susceptible to oxidative alteration due to greater 4-hydroxynonenal (4-HNE) concentrations per unit of intramuscular triglycerides in obese individuals. Obese individuals may also have a larger target for ROS oxidation because of the higher amounts of lipid molecules in their bodies. Excessive fat builds up in WAT was found to enhance lipid peroxidation within the WAT itself, according to research using animal models of obesity by Furukawa and colleagues. A decrease in the activity of antioxidant enzymes such as Superoxide Dismutase (SOD), catalase (CAT), and Glutathione Peroxidase (GPx) in WAT was found in animal experiments to be linked to obesity. The consumption of lipids in the diet also contributes to the development of chronic oxidative stress in the body. In middle-aged men with abdominal obesity, conjugated linolenic acid consumption elevated urine concentrations of 8-epi PGF_{2a}.³⁵⁻³⁹

CONCLUSION

It is concluded from the above discussion and review *Ama* is the first and main culprit. This causes the *Sthaulya*. There is also a clear imbalance between *Ama* formations and their removal or

neutralization; in terms of free radicals, we can say that free radical load is more, and antioxidant defence is not enough to remove these free radicals. It can be either due to excessive free radical formation or a weakened antioxidant defence system leading to Oxidative stress. It is also concluded from the above discussion that *Ama* disrupts the *Dhatu Poshan Nayayas* and creates *Sama Dhatus*, and maximum disruption is at the *Meda Dhatu*. The lowest antioxidant activity occurs in the adipose tissue, causing high oxidative stress and leading to inflammatory and pro-inflammatory states of hyperplasia and hypertrophy. This explains the complex pathology of *Sthaulya* disease.

REFERENCES

- Zhang ZY, Wang MW. Obesity, a health burden of a global nature. *Acta Pharmacol Sin* 2012;33:145–147.
- Pollin IS, Kral BG, Shattuck T, et al. High prevalence of cardiometabolic risk factors in women considered low risk by traditional risk assessment. *J Womens Health (Larchmt)* 2008;17:947–953.
- Jung RT. Obesity as a disease. *Br Med Bull* 1997;53: 307–321.
- Pi-Sunyer FX. Health implications of obesity. *Am J Clin Nutr* 1991;53:1595S–1603S.
- Sikaris KA. The clinical biochemistry of obesity. *Clin Biochem Rev* 2004;25:165–181.
- Despres JP, Moorjani S, Lupien PJ, et al. Regional distribution of body fat, plasma lipoproteins, and cardiovascular disease. *Arteriosclerosis* 1990;10:497–511.
- Fernandez-Sanchez A, Madrigal-Santillan E, Bautista M, et al. Inflammation, oxidative stress, and obesity. *Int J MolSci* 2011;12:3117–3132.
- Savini I, Catani MV, Evangelista D, et al. Obesity associated oxidative stress: Strategies finalized to improve redox state. *Int J Mol Sci* 2013;14:10497–10538.
- Furukawa S, Fujita T, Shimabukuro M, et al. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest* 2004;114:1752–1761.
- Lee H, Lee YJ, Choi H, et al. Reactive oxygen species facilitate adipocyte differentiation by accelerating mitotic clonal expansion. *J Biol Chem* 2009;284:10601–10609.
- Higuchi M, Dusing GJ, Peshavariya H, et al. Differentiation of human adipose-derived stem cells into fat involves reactive oxygen species and Forkhead box O1 mediated upregulation of antioxidant enzymes. *Stem Cells Dev* 2013;22:878–888.
- Horvath TL, Andrews ZB, Diano S. Fuel utilization by hypothalamic neurons: Roles for ROS. *Trends Endocrinol Metab* 2009;20:78–87.
- Serra D, Mera P, Malandrino MI, et al. Mitochondrial fatty acid oxidation in obesity. *Antioxid Redox Signal* 2013;19:269–284.
- Beltowski J. Leptin and the regulation of endothelial function in physiological and pathological conditions. *Clin Exp Pharmacol Physiol* 2012;39:168–178.
- Chrysohoou C, Panagiotakos DB, Pitsavos C, et al. The implication of obesity on total antioxidant capacity in apparently healthy men and women: The ATTICA study. *Nutr Metab Cardiovasc Dis* 2007;17:590–597.
- Patel C, Ghanim H, Ravishankar S, et al. Prolonged reactive oxygen species generation and nuclear factor kappa B activation after a high-fat, high-carbohydrate meal in the obese. *J Clin Endocrinol Metab* 2007;92:4476–4479.
- Neha Rawat and Rakesh et al. Correlation of Rasa dhatu in Modern Perspective: A Review. *Int. J. Res. Ayurveda Pharm.* 2019; 10(2):28-32 <http://dx.doi.org/10.7897/2277-4343.100231>
- Charaka Samhita (text with English translation & critical Exposition based on Chakrapani Dutta Ayurveda Dipika) Vol IV (Chikitsa sthana 15/16) edited by R. K Sharma and Bhagwan Das, Chaukhamba Sanskrit series Office Varanasi (India), Reprint 2016.
- Illustrated Sushruta Samhita (Text, English Translation. Notes, Appendices, and Index) Vol 1 (sutra sthana14/10) Translator Prof. K.R Srikantha Murthy, Chaukhamba Orientalia, Varanasi, reprint 2016.
- Charaka Samhita (text with English translation & critical Exposition based on Chakrapani Dutta Ayurveda Dipika) Vol 1 (sutra sthana 28/4) edited by R. K Sharma and Bhagwan das, Chaukhamba Sanskrit series Office Varanasi (India), Reprint 2016.
- Illustrated Sushruta Samhita (Text, English Translation. Notes, Appendices, and Index) Vol 1 (sutra sthana 46/528) Translator Prof. K.R Srikantha Murthy, Chaukhamba Orientalia, Varanasi, reprint 2016.
- Illustrated Sushruta Samhita (Text, English Translation. Notes, Appendices, and Index) Vol 1 (sutra sthana14/14) Translator Prof. K.R Srikantha Murthy, Chaukhamba Orientalia, Varanasi, reprint 2016.
- Ortega RM, Rodriguez-Rodriguez E, Aparicio A, et al. Young children with excess of weight show an impaired selenium status. *Int J Vitam Nutr Res* 2012;82:121–129.
- Strauss RS. Comparison of serum concentrations of alpha-tocopherol and beta-carotene in a cross-sectional sample of obese and nonobese children (NHANES III). *National Health and Nutrition Examination Survey. J Pediatr* 1999; 134:160–165.
- Schleicher RL, Carroll MD, Ford ES, et al. Serum vitamin C and the prevalence of vitamin C deficiency in the United States: 2003–2004 National Health and Nutrition Examination Survey (NHANES). *Am J Clin Nutr* 2009;90:1252–1263.
- Reitman A, Friedrich I, Ben-Amotz A, et al. Low plasma antioxidants and normal plasma B vitamins and homocysteine in patients with severe obesity. *Isr Med Assoc J* 2002;4:590–593.
- Kaidar-Person O, Person B, Szomstein S, et al. Nutritional deficiencies in morbidly obese patients: A new form of malnutrition? Part A: Vitamins. *Obes Surg* 2008;18: 870–876.
- Andersen LF, Jacobs DR, Jr., Gross MD, et al. Longitudinal associations between body mass index and serum carotenoids: The CARDIA study. *Br J Nutr* 2006;95:358–365.
- Kuno T, Hozumi M, Morinobu T, et al. Antioxidant vitamin levels in plasma and low density lipoprotein of obese girls. *Free Radic Res* 1998;28:81–86.
- Aasheim ET, Bohmer T. Low preoperative vitamin levels in morbidly obese patients: A role of systemic inflammation? *Surg Obes Relat Dis* 2008;4:779–780.
- Canoy D, Wareham N, Welch A, et al. Plasma ascorbic acid concentrations and fat distribution in 19,068 British men and women in the European Prospective Investigation into Cancer and Nutrition Norfolk cohort study. *Am J Clin Nutr* 2005;82:1203–1209.
- Myara I, Alamowitch C, Michel O, et al. Lipoprotein oxidation and plasma vitamin E in nondiabetic normotensive obese patients. *Obes Res* 2003;11:112–120.
- Weisstaub G, Hertrampf E, Lopez de Romana D, et al. Plasma zinc concentration, body composition and physical activity in obese preschool children. *Biol Trace Elem Res* 2007;118:167–174.
- Kaidar-Person O, Person B, Szomstein S, et al. Nutritional deficiencies in morbidly obese patients: A new form of malnutrition? Part B: Minerals. *Obes Surg* 2008;18: 1028–1034.

35. Bakker SJ, IJzerman RG, Teerlink T, *et al.* Cytosolic triglycerides and oxidative stress in central obesity: The missing link between excessive atherosclerosis, endothelial dysfunction, and beta-cell failure? *Atherosclerosis* 2000;148:17–21.
36. Inoguchi T, Li P, Umeda F, *et al.* High glucose level and free fatty acid stimulate reactive oxygen species production through protein kinase C—dependent activation of NAD(P)H oxidase in cultured vascular cells. *Diabetes* 2000;49:1939–1945.
37. Russell AP, Gastaldi G, Bobbioni-Harsch E, *et al.* Lipid peroxidation in skeletal muscle of obese as compared to endurance-trained humans: A case of good vs bad lipids? *FEBS Lett* 2003;551:104–106.
38. Vincent HK, Powers SK, Dirks AJ, *et al.* Mechanism for obesity-induced increase in myocardial lipid peroxidation. *Int J Obes Relat Metab Disord* 2001;25:378–388.
39. Basu S, Riserus U, Turpeinen A, *et al.* Conjugated linoleic acid induces lipid peroxidation in men with abdominal obesity. *Clin Sci (Lond)* 2000;99:511–516.

Cite this article as:

Amulya Murthy Aku *et al.* Sthaulya, ama, and meda dhatu, The dysfunctional nutritional metabolism with special reference to oxidative stress: A narrative review. *Int. J. Res. Ayurveda Pharm.* 2022;13(4):95-98 <http://dx.doi.org/10.7897/2277-4343.130496>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IJRAP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publishing quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJRAP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IJRAP editor or editorial board members.